

Cancer: Complementary/Alternative Medical Care

Carotene (Vitamin A Precursor) 500,000 iu/d

Niacin (B3) 1 gm tid (Adequate Vitamin C helps prevent flush).

Inositol hexanicotinate prevents flush/metabolized to B3, Niacinamide also works at this dose

Vitamin C Highest tolerated dose

<http://www.doctoryourself.com/cameron.html>

Vitamin E 800-1200 iu/d

B Complex 100 mg/d

Selenium 200-1,000 ug/d

Ca++ 500 mg tid

Mg++ 250 mg tid

<http://www.doctoryourself.com/gersonspeech.html>

high potassium, low sodium diet, with no fats or oils, and minimal animal proteins. Juices of raw fruits and vegetables and of raw liver provide active oxidizing enzymes, which facilitate rehabilitation of the liver. Iodine and niacin supplementation is used. Caffeine enemas cause dilation of bile ducts, which facilitates excretion of toxic cancer breakdown products by the liver and dialysis of toxic products from blood across the colonic wall.

Folic acid 1-30 mg/d

CoQ10 300-600 mg/d (esp. breast CA)

Chlorophyll

Bladder Cancer B6

Lung Cancer Folate

Zn Prostate CA (40-150 mg/d)

Cu Inverse relationship with Zn